



# Computational Thinking

## Solutions to Exercise 6 (Cryptography)

### 1 Zero Knowledge Proofs in Geometry

- a) The constructions are simple and we show here for example how to bisect an angle. First we open the compass in an arbitrary angle and draw a circle around the endpoint of the angle. We label the intersection points of the circle and the angle as  $A$  and  $B$ . We draw a circle around the point of  $A$  and  $B$ . And we construct a line between the endpoint of the angle and the intersection of the (newly constructed) circles <sup>1</sup>.
- b) The following example is one of the possible protocols:

#### ZKP in Geometry

Peggy		Vic
knows $\alpha, \beta = 3\alpha$		knows $\beta$
create random angle $\gamma$		
construct $\tau = 3\gamma$	$\xrightarrow{\text{send over } \tau}$	
	$\xleftarrow{\text{send over } c}$	choose randomly $c \in \{0, 1\}$
create $\rho = \gamma + c\alpha$	$\xrightarrow{\text{send over } \rho}$	check $3\rho \stackrel{?}{=} \tau + c\beta$

- **Completeness.** One can easily see that if Peggy is honest and knows  $\alpha$ , Vic always accepts. More concretely, in the last step  $3\rho = 3(\gamma + c\alpha) = 3\gamma + 3c\alpha = \tau + c\beta$ .
- **Soundness.** We show that if Peggy can answer both challenges then she really knows  $\alpha$ . Assume Peggy can answer for both challenges  $c = 0$  and  $c' = 1$  correctly with  $\rho = \gamma + 0 * \alpha = \gamma$  and  $\rho' = \gamma + \alpha$ . Then it follows that Peggy can compute  $\alpha = \rho' - \rho$ . In other words, if she doesn't know  $\alpha$  she can at most answer one of the challenges, and fail at the other challenge. That is, Peggy can correctly answer in one round only with probability  $1/2$ , and therefore  $n$  rounds only with probability  $1/2^n$ .
- **Zero Knowledge.** The main idea is to show that the same<sup>2</sup> transcript that Victor has after the protocol could be generated by himself (without knowing  $\alpha$ ). During the protocol the transcript contains the triples  $(\tau, c, \rho)$  and can be produced as follows. For each challenge  $c$ , generate a random  $\rho$  and construct  $\tau = 3\rho - c\beta$ .

<sup>1</sup>You may have a look at the following video about these constructions and the impossibility of trisecting an angle, if desired: <https://youtu.be/01sPvUr0YCO>


<sup>2</sup>With the same distribution, in case of random values.

*Note.* The soundness and zero knowledge property may sound contradictory to each other but they are not. Even though the transcript has "no information about  $\alpha$ " after the protocol, Victor is convinced that Peggy knows  $\alpha$  because she was able to answer to the challenges *during* the protocol. In particular, first  $\tau$  is constructed and *only then* based on the challenge  $c$ ,  $\rho$  is constructed.

## 2 MPC with Secret Sharing

a) One can easily see that each party after summing locally holds a share of the polynomial  $f(x) = f_1(x) + f_2(x)$  (since the degree of the polynomial by summing doesn't change and  $t$  points uniquely define a polynomial of degree  $t - 1$ ). It follows,  $s = f(0) = f_1(0) + f_2(0) = s_1 + s_2$ . Hence, if the polynomial is reconstructed and evaluated at point 0, it will result to the sum of  $s_1$  and  $s_2$ .

b) Dave can just continue the same way as other participants:



c) Alice and Carol can easily compute the salary of Bob as follows. Since Carol has  $m_2 = b + m_1$  and Alice has  $m_1$  they can compute  $b = m_2 - m_1$ . The same technique can be used for Dave as well, namely  $d = m_4 - m_3$ .

d) The main idea is to use secret sharing and the fact that they are linear. Each party shares their salary by using  $(n, n)$  Shamir secret sharing, compute locally the sum of each share (namely the share of each salary), and in the end reconstruct the sum only.