


Discrete Event Systems


Exercise Sheet 3

1 From DFA to Regular Expression


First generate the GNFA.




Then remove node 2.



Then remove node 3.




Then remove node 1 and derive the corresponding regular expression.




Note that we could also start by removing the node 3.

2 Transforming Automata [Exam HS14]

The regular expression can be obtained from the finite automaton using the transformation presented in the script. After ripping out state q_2 , the corresponding GNFA looks like this:



After also removing state q_1 , the GNFA looks as follows.



Eliminating the last state q_3 yields the final solution, which is $(01^*0)^*1(0 \cup 11^*0(01^*0)^*1)^*$.

Note: Ripping out the interior states in a different order yields a distinct yet equivalent regular expression. The order q_3, q_2, q_1 , for example, results in $((0 \cup 10^*1)1^*0)^*10^*$.

3 Pumping Lemma

The Pumping Lemma in a Nutshell

Given a language L , assume for contradiction that L is regular and has the pumping length p . Construct a suitable word $w \in L$ with $|w| \geq p$ (“there exists $w \in L$ ”) and show that for all divisions of w into three parts, $w = xyz$, with $|x| \geq 0$, $|y| \geq 1$, and $|xy| \leq p$, there exists a pumping exponent $i \geq 0$ such that $w' = xy^iz \notin L$. If this is the case, L is not regular.

a) We claim that L_1 is not regular and prove our claim with the pumping lemma recipe:

1. Assume for contradiction that L_1 was regular.
2. There must exist some p , s.t. any word $w \in L_1$ with $|w| \geq p$ is pumpable.
3. Choose the string $w = 1^p02^p \in L_1$ with length $|w| > p$.
4. Consider all ways to split $w = xyz$ s.t. $|xy| \leq p$ and $|y| \geq 1$.
→ Hence, $y \in 1^+$.
5. Observe that $xy^0z \notin L_1$ – a contradiction to p being a valid pumping length.
6. Consequently, L_1 cannot be regular.

b) Language L_2 can be shown to be non-regular using the pumping lemma. Assume for contradiction that L_1 is regular and let p be the corresponding pumping length. Choose w to be the word 0110^p1^p . Because w is an element of L_1 and has length more than p , the pumping lemma guarantees that w can be split into three parts, $w = xyz$, where $|xy| \leq p$ and for any $i \geq 0$, we have $xy^iz \in L_1$. In order to obtain the contradiction, we must prove that for every possible partition into three parts $w = xyz$ where $|xy| \leq p$, the word w cannot be pumped. We therefore consider the various cases.

- (1) If y starts anywhere within the first three symbols (i.e. 011) of w , deleting y (pumping with $i = 0$) creates a word with an illegal prefix (e.g. $10^p 1^p$ for $y = 01$).
- (2) If y consists of only 0s from the second block, the word $w' = xy^2z$ has more 0s than 1s in the last $|w'| - 3$ symbols and hence $c \neq d$.

Note that y cannot contain 1s from the second block because of the requirement $|xy| \leq p$.

We have shown that for all possible divisions of w into three parts, the pumped word is not in L_1 . Therefore, L_1 cannot be regular and we have a contradiction.

Note that we could have also used the pumping lemma recipe to prove that L_2 is not regular.

Be Careful!

The argumentation above is based on the closure properties of regular languages and only works in the direction presented. That is, for an operator $\diamond \in \{\cup, \cap, \bullet\}$, we have:

If L_1 and L_2 are regular, then $L = L_1 \diamond L_2$ is also regular.

If either L_1 or L_2 or both are non-regular, we cannot deduce the non-regularity of L or vice-versa. Moreover, L being regular does not imply that L_1 and L_2 are regular as well. This may sound counter-intuitive which is why we give examples for the three operators.

- $L = L_1 \cup L_2$: Let L_1 be any non-regular language and L_2 its complement. Then $L = \Sigma^*$ is regular.
- $L = L_1 \cap L_2$: Let L_1 be any non-regular language and L_2 its complement. Then $L = \emptyset$ is regular.
- $L = L_1 \bullet L_2$: Let $L_1 = \{a^*\}$ (a regular language) and $L_2 = \{a^p \mid p \text{ is prime}\}$ (a non-regular language) then $L = \{aaa^*\}$ is regular.

Hence, to prove that a language L_x is non-regular, you assume it to be regular for contradiction. Then you combine it with a *regular* language L_r to obtain a language $L = L_x \diamond L_r$. If L is non-regular, L_x could not have been regular either.

4 Pumping Lemma Revisited

- a) Let us assume that L is regular and show that this results in a contradiction.

We have seen that any regular language fulfills the pumping lemma. This means, there exists a number p , such that every word $w \in L$ with $|w| \geq p$ can be written as $w = xyz$ with $|xy| \leq p$ and $|y| \geq 1$, such that $xy^i z \in L$ for all $i \geq 0$.

In order to obtain the contradiction, we need to find at least one word $w \in L$ with $|w| \geq p$ that does not adhere to the above proposition. We choose $w = xyz = 1^{p^2}$ and consider the case $i = 2$ for which the Pumping Lemma claims $w' = xy^2z \in L$.

We can relate the lengths of $w = xyz$ and $w' = xy^2z$ as follows.

$$p^2 = |w| = |xyz| < |w'| = |xy^2z| \leq p^2 + p < p^2 + 2p + 1 = (p + 1)^2$$

So we have $p^2 < |w'| < (p + 1)^2$ which implies that $|w'|$ cannot be a square number since it lies between two consecutive square numbers. Therefore, $w' \notin L$ and hence, L cannot be regular.

- b) Consider the alphabet $\Sigma = \{a_1, a_2, \dots, a_n\}$ and the language $L = \bigcup_{i=1}^n a_i^* = a_1^* \cup a_2^* \cup \dots \cup a_n^*$. In other words, each word of the language L contains an arbitrary number of just **one** symbol a_i . The language is regular, as it is the union of regular languages, and the smallest possible pumping number p for L is 1. But any DFA needs at least $n + 2$ states to accept

the empty word, distinguish the n different characters of the alphabet, and for a failing state. Thus, for the DFA, we cannot deduce any information from p about the minimum number of states.

The same argument holds for the NFA.

5 Minimum Pumping Length

To begin with, observe that the minimum pumping length p of a language $L = L_1 \cup L_2$ is at most $p \leq \max\{p_1, p_2\}$, where p_1 and p_2 are the minimum pumping lengths of L_1 and L_2 , respectively. This holds because if there is already a string w that is pumpable in L_1 , then w will also be pumpable in L . Hence, let $L_1 = 1^*0^+1^+0^*$ and $L_2 = 111^+0^+$.

- The minimum pumping length of L_2 cannot be 4 because 1110 cannot be pumped. Now consider the string s that belongs to L_2 and that has a size of 5. If $s = 11110$, then it can be divided into xyz where $x = 111$, $y = 1$ and $z = 0$ and thus can be pumped. If $s = 11100$, then it can be divided into xyz where $x = 111$, $y = 0$ and $z = 0$ and thus can be pumped. Similarly, all longer words can be pumped. The minimum pumping length for L_2 is thus 5.
- A string s of size 3 and belonging to L_1 can always be pumped.

Considering the word 1110, observe that it can also not be pumped in $L = L_1 \cup L_2$. In conclusion, the minimum pumping length of L is 5.

6 The art of being regular

L is not regular. We show it using the pumping lemma. We start by choosing a string in L . Let $w = 100^p\#10^p$. Then $w \in L$ since $x(100^p)$ is equal to $2y$ (where y is 10^p) for $p \geq 0$. We must consider three cases for where y can fall:

- $y = 1$ Pump out. Arithmetic is wrong. The left side is 0 but right side isn't.
- $y = 10^*$ Pump out. Arithmetic is wrong.
- $y = 0^p$ Pump out. Arithmetic is wrong. Decreased left side but not right. So, in particular, it is no longer the case that $x \geq y$ (required since $y \neq \emptyset$).

Bonus tasks: – solutions provided by student Angéline Pouget in HS20

- Determine whether $L = \{x\#y \mid x + y = 3y\}$ is context-free.

To begin with, we observe that

$$\begin{aligned} L &= \{x\#y \mid x + y = 3y\} \\ &= \{x\#y \mid x = 2y\} \\ &= \{w0\#w \mid w \in 1(0 \cup 1)^*\}. \end{aligned}$$

We prove that $L = \{w0\#w \mid w \in 1(0 \cup 1)^*\}$ is not context-free using the tandem-pumping lemma. First, we assume for contradiction that L is context-free and hence there is a number p such that any string in L of length $\geq p$ is tandem-pumpable within a substring of length p . We choose $w = 1^p0^p$ and thereby consider the word $\alpha = w0\#w = 1^p0^p0\#1^p0^p$ with $|\alpha| \geq p$.

We now want to split $\alpha = uvxyz$ with $|vy| \geq 1$, $|vxy| \leq p$ and $w^i xy^i z \in L$ for all $i \geq 0$. Because we have $|vxy| \leq p$, there are the following options:

- $\# \notin vxy$ ($vxy = 1^m$ or $vxy = 0^m$ with $1 \leq m \leq p$ or $vxy = 1^n0^s$ with $n + s \leq p$). Any one of these sequences can either be before or after the $\#$ but independent of this choice, if we pump v and y and choose for example $i = 0$, we will have $\alpha' = w'0\#w''$ with $w' \neq w$ and hence $\alpha' \notin L$.

- $\# \in vxy$. In this case, we can choose $x = \#$ because we know that there is only one $\#$ and therefore this cannot be the pumpable part. This leaves us with $v = 0^n$ and $y = 1^s$ with $1 \leq n + s \leq p - 1$ and if we for example set $i = 0$ this leaves us with $\alpha' = 1^p 0^{p+1-n} \# 1^{p-s} 0^p$ which is $\notin L$.


Because we have now considered all possible splits of this word into $\alpha = uvxyz$, we can safely say that language L is not context-free.

- Show whether $L' = \{x\#y \mid x + \text{reverse}(y) = 3 \cdot \text{reverse}(y)\}$ is context-free. The $\text{reverse}()$ -function takes an integer as a bitstring and reverses the order of its bits.

Let $w' = \text{reverse}(w)$. Applying the same transformations as above, we obtain

$$L' = \{x\#y \mid x = 2 \cdot \text{reverse}(y)\} = \{w0\#w' \mid w \in 1(0 \cup 1)^*\}.$$

We can show that this language is context-free by drawing a push-down automaton that accepts this language. This automaton is depicted below with “>” representing stack operations “→”.



We could have alternatively shown that the language is context-free by providing a context free grammar (V, Σ, R, S) such as the following:

- $V = \{S\}$
- $\Sigma = \{0, 1, \#\}$
- $R: S \rightarrow 1S1 \mid 0S0 \mid 0\#$
- $S = S$